

TWO STUDIES IN CENTRAL ASIAN SEISMOLOGY: A TELESEISMIC STUDY OF
THE PAMIR/HINDU KUSH SEISMIC ZONE AND ANALYSIS OF DATA FROM
THE KYRGYZSTAN BROADBAND SEISMIC NETWORK

Robert John Mellors

Submitted to the faculty of the Graduate School

in partial fulfillment of the

requirements for the degree

Doctor of Philosophy

in the Department of Geological Sciences

Indiana University

June, 1995

copyright, 1995

Robert John Mellors

ALL RIGHTS RESERVED

In memory of the victims of the Suusamyр earthquake.

It is my hope that perhaps this work may, in some small manner,
help alleviate the suffering from future catastrophes.

Acknowledgments

To thank everyone who has aided me in this work would require several gigabytes of disk space. Unfortunately, I must attempt to compress it into one page.

I would like to thank Dr. Michael Hamburger for his patience as an advisor. His insights on both seismology and Central Asia were invaluable and gave direction to my work. I would also like to thank Dr. Gary Pavlis for his advice and enthusiasm for all things seismological and especially for his efforts in creating the hacker's paradise of equipment and programs we have in the geophysics lab. The efforts of Dr. Frank Vernon built the foundations on which much of this work is constructed. He deserved thanks for allowing me to work on the KNET network and his patience in explaining the basics of the KNET network, sometimes in trying conditions, will always be appreciated.

I also would like to thank Dr. Albert Rudman, Dr. Bruce Douglas, and Dr. Peter Sternberg, for their reading of this manuscript and helpful discussions on topics ranging from Laramide thrusting to synthetic seismograms and partial differential equations.

The North Lab, especially Glen Offield, Phil Porter, and Marina Glushko deserve thanks both for their technical contributions and hard work but for also their unfailing support (sometimes literally) and friendship during many months of rigorous field work both in Central Asia and La Jolla.

Our many colleagues in Kyrgyzstan provided much help during our work there and this work could not have been done without them. Vitali Bragin, Shumkarbek Jumadenov, Dulat Raskulov, Larisa Rudakova, and Sasha Zubovich were particularly helpful. I also thank all the station personnel for their hospitality.

This work was greatly speeded by the elegant and efficient software developed by Danny Harvey and Dan Quinlan at the Joint Seismic Center at the University of Colorado. David Simpson graciously provided the digital topographic data which has been invaluable for many of the figures in this work. Discussions with Vitaly Khalturin,

Tatiana Rautian, Vladik Martynov, Sujoy Ghose, Andrei Korjenkov, Haydar Al-Shukri and Steve Roecker helped provoke, develop and refine the ideas in this thesis.

I am also extremely appreciative of the financial support from the Joint Seismic Program of the Incorporated Research Institutions in Seismology (IRIS) during my work here.

I especially would like to thank my parents for their support through all these years.

During my years in Bloomington (and Bishkek) I had the support of many friends and a big "Beh!" goes to all of them: Paul Anderson, Glen and Lorie Bear, Steve Bennett, Julie Boyd, John Hohman, the softball gang, Priya Ganguli, Pete Graydon, Gary Rockow, Steven King, Irina Kotozhekova, Dave Millen, Sue McDonald, Manoling Ramos, and Stephanie Werne.

Finally, I would like to thank Oreo the cat for her efforts in keeping mice away from my thesis.

Chong

ragmad

y'all!

Two studies in Central Asian seismology: A teleseismic study of the Pamir/Hindu Kush seismic zone and analysis of data from the Kyrgyzstan broadband seismic network

Robert J. Mellors

The thesis consists of two separate studies: (1) an analysis of lateral velocity variations associated with the Pamir and Hindu Kush seismic zones and (2) a study of the seismicity and tectonics of the Kyrgyz Tien Shan, with emphasis on a major earthquake and its aftershock sequence. In the first study, I used teleseismic arrival time data from analog seismic stations in Tadjikistan and northern Afghanistan to invert for velocity perturbations in the crust and upper mantle of the Pamir-Hindu Kush. The results show a strong (up to 6%) well-resolved zone of high velocities in the upper mantle at depths greater than 200 km, coincident with the location of the deep Hindu Kush earthquakes. No clear velocity perturbations are associated with the deep Pamir earthquakes. Above 200 km little correlation is observed between the velocity perturbations and the seismic zone, but indications of thicker crust under the Pamir and thinner crust under the Tadjik Depression are seen. I interpret these high velocities under the Hindu Kush as evidence of oceanic lithosphere at depth, possibly resulting from the subduction of an isolated section of oceanic lithosphere. The lack of velocity anomalies in the upper segments of the Hindu Kush and Pamir seismic zones may indicate the subduction of continental lithosphere.

The second study concentrated on data from the Kyrgyzstan broadband seismic network (KNET), which consists of 10 telemetered broadband seismometers installed in the area around Bishkek, Kyrgyzstan. In 3 1/2 years of operation it has recorded two nearby aftershock sequences, and numerous local, regional, and teleseismic events. Most local earthquakes are concentrated along the Kyrgyz Range in a diffuse pattern, suggesting that the seismicity occurs throughout a complex set of faults rather than on

one main range-bounding fault. Many quarry blasts were also recorded, which allowed calibration of velocity models for network locations. The high dynamic range and broadband response of the KNET seismometers permits observation of a range of phenomena seldom observed by conventional systems. These include: (1) a ramp-like phase between the P and the S wave arrivals that is observed on several dozen nearby (< 15 km distance) events. This ramp probably represents the effect of near-field source terms; (2) a pronounced vertical displacement signal that coincides with the arrival of the S wave for a number of high-amplitude local events. This displacement appears to be due to an instrumental or site effect; (3) strong variations in aftershock waveforms that appear to be dependent in part on source depth; and (4) variations in S wave amplitude across the network from regional events, especially those from the south.

The Ms 7.4 Suusamyr, Kyrgyzstan earthquake of 19 August 1992 was studied using aftershock data, a teleseismic body wave inversion, and field observations. The combined analysis defined a 50 km-long, east-west rupture zone that dips 50° to the south and extends to a depth of 18 km. The rupture originated at the eastern end at depth and propagated westward, and probably included at least one sub-event. The three large (Ms > 6.0) aftershocks that occurred immediately after the main shock also showed a progression with time from east to west. A strong correlation between the aftershock zone and topography was noted, suggesting that the topography is largely fault controlled. The Suusamyr event demonstrates that a large amount of current shortening is accommodated along high-angle reverse faults within the Tien Shan in addition to that occurring along the range-bounding fault systems. This event also clearly shows the 3-dimensional geometry of an active, basement-involved, intra-continental thrust.

ACKNOWLEDGMENTS.....	4
LIST OF FIGURES	10
CHAPTER 1:	13
INTRODUCTION	13
REFERENCES.....	15
CHAPTER 2:	ERROR! BOOKMARK NOT DEFINED.
EVIDENCE FOR A HIGH-VELOCITY SLAB ASSOCIATED WITH THE HINDU KUSH SEISMIC ZONE	ERROR! BOOKMARK NOT DEFINED.
ABSTRACT	ERROR! BOOKMARK NOT DEFINED.
INTRODUCTION	ERROR! BOOKMARK NOT DEFINED.
TECTONIC SETTING.....	ERROR! BOOKMARK NOT DEFINED.
DATA.....	ERROR! BOOKMARK NOT DEFINED.
<i>Collection</i>	<i>Error! Bookmark not defined.</i>
<i>Reduction</i>	<i>Error! Bookmark not defined.</i>
TELESEISMIC INVERSION.....	ERROR! BOOKMARK NOT DEFINED.
PROCEDURE.....	ERROR! BOOKMARK NOT DEFINED.
RESULTS.....	ERROR! BOOKMARK NOT DEFINED.
<i>Layer 1: Crustal layer</i>	<i>Error! Bookmark not defined.</i>
<i>Layer 2: Mantle (50-200 km)</i>	<i>Error! Bookmark not defined.</i>
<i>Layer 3: Mantle (200-350 km)</i>	<i>Error! Bookmark not defined.</i>
<i>Error and resolution</i>	<i>Error! Bookmark not defined.</i>
DISCUSSION AND CONCLUSIONS.....	ERROR! BOOKMARK NOT DEFINED.
REFERENCES.....	ERROR! BOOKMARK NOT DEFINED.
CHAPTER 3:	ERROR! BOOKMARK NOT DEFINED.
THE KYRGYZSTAN BROADBAND SEISMIC NETWORK: DESCRIPTION, SETTING, AND PRELIMINARY RESULTS	ERROR! BOOKMARK NOT DEFINED.
INTRODUCTION.....	ERROR! BOOKMARK NOT DEFINED.
GEOLOGIC AND TECTONIC SETTING.....	ERROR! BOOKMARK NOT DEFINED.
<i>Crustal and upper mantle structure</i>	<i>Error! Bookmark not defined.</i>
<i>Tectonics</i>	<i>Error! Bookmark not defined.</i>
<i>Historical seismicity</i>	<i>Error! Bookmark not defined.</i>
<i>Instrumentally recorded seismicity</i>	<i>Error! Bookmark not defined.</i>
TECHNICAL DESCRIPTION.....	ERROR! BOOKMARK NOT DEFINED.
PRELIMINARY RESULTS.....	ERROR! BOOKMARK NOT DEFINED.
<i>Velocity models and comparison with KNET data</i>	<i>Error! Bookmark not defined.</i>
<i>Local seismicity</i>	<i>Error! Bookmark not defined.</i>
<i>Waveform observations</i>	<i>Error! Bookmark not defined.</i>
<i>Near-field phases</i>	<i>Error! Bookmark not defined.</i>
<i>Vertical offsets</i>	<i>Error! Bookmark not defined.</i>
<i>Regional events</i>	<i>Error! Bookmark not defined.</i>
CONCLUSIONS AND SUGGESTIONS FOR FUTURE WORK.....	ERROR! BOOKMARK NOT DEFINED.
REFERENCES.....	ERROR! BOOKMARK NOT DEFINED.
CHAPTER 4:	ERROR! BOOKMARK NOT DEFINED.
THE 1992 SUUSAMYR, KYRGYZSTAN, EARTHQUAKE	ERROR! BOOKMARK NOT DEFINED.
ABSTRACT	ERROR! BOOKMARK NOT DEFINED.
INTRODUCTION.....	ERROR! BOOKMARK NOT DEFINED.
<i>Geologic and tectonic setting</i>	<i>Error! Bookmark not defined.</i>
<i>Field observations</i>	<i>Error! Bookmark not defined.</i>

<i>Main shock</i>	<i>Error! Bookmark not defined.</i>
AFTERSHOCK DATA.....	ERROR! BOOKMARK NOT DEFINED.
<i>Data and processing</i>	<i>Error! Bookmark not defined.</i>
<i>Results</i>	<i>Error! Bookmark not defined.</i>
BODY WAVE INVERSION	ERROR! BOOKMARK NOT DEFINED.
DISCUSSION AND CONCLUSIONS.....	ERROR! BOOKMARK NOT DEFINED.
<i>Comparison between aftershock and inversion results</i>	<i>Error! Bookmark not defined.</i>
<i>Aftershock locations and geometry of the fault plane</i>	<i>Error! Bookmark not defined.</i>
<i>Lack of surface rupture.</i>	<i>Error! Bookmark not defined.</i>
<i>Regional tectonics.</i>	<i>Error! Bookmark not defined.</i>
<i>Seismic hazard</i>	<i>Error! Bookmark not defined.</i>
REFERENCES.....	ERROR! BOOKMARK NOT DEFINED.
CONCLUSIONS.....	ERROR! BOOKMARK NOT DEFINED.
APPENDIX A:	ERROR! BOOKMARK NOT DEFINED.
STATION LOCATIONS AND DESCRIPTIONS	ERROR! BOOKMARK NOT DEFINED.
APPENDIX A:	ERROR! BOOKMARK NOT DEFINED.
INSTITUTE OF PHYSICS OF THE EARTH TEMPORARY AFTERSHOCK NETWORK DATA PROCESSING	ERROR! BOOKMARK NOT DEFINED.

List of figures

- Chapter 1 Introduction
 - 1.1 Tectonic map of Asia
- Chapter 2 Evidence for a high-velocity slab associated with the Hindu Kush seismic zone
 - 2.1 Regional map of Hindu Kush seismic zone
 - 2.2 Earthquakes recorded by the Soviet regional network
 - 2.3 Depth to Moho
 - 2.4a Teleseismic earthquakes
 - 2.5a Relative residuals and map
 - 2.5b Relative residuals
 - 2.6 Averaged results of inversion
 - 2.7 Single inversion results
 - 2.8 Resolution matrix of results
 - 2.9 Cross section of results and earthquakes
- Chapter 3 The KNET network
 - 3.1 Map of Kyrgyzstan broadband network
 - 3.2 Simplified diagram of data flow in the KNET network
 - 3.3 Schematic of typical station hardware configuration
 - 3.4 Photo of typical IVTAN mountain station (top)
and KIS valley station (bottom)
 - 3.5 Photo of STS-2 seismometer before installation (top) and photo of inside of vault (bottom).
 - 3.6 Phase and amplitude responses for Streckeisen STS-2 sensors with anti-aliasing filters for 100 sps data stream.

- 3.7 Seismogram of local earthquake as recorded by 20 sps and 100 sps data stream and showing 'ringing' effect of anti-aliasing filters.
- 3.8 Map of northern Tien Shan with CMT solutions and large historical earthquakes.
- 3.9 Map of regional seismicity in northern Tien Shan between 1929 and 1950
- 3.10 Regional seismicity according to the Soviet regional catalog.
- 3.11 Local seismicity recorded by KNET between 1991 and 1995 showing Suusamyr aftershocks and quarry blasts.
- 3.12 Velocity models for the northern Tien Shan.
- 3.13 Comparison of quarry blast P travel times with theoretical arrival times predicted by velocity models (reduced by 7km/s)
- 3.3.4a Quarry blast seismogram showing air wave (top) and diagram of air wave arrival time plotted versus P - S time.
- 3.15 Seismicity maps and cross sections of the Bishkek region
- 3.16 Focal mechanisms in the Bishkek region
- 3.17 Seismogram of local event near station USP.
- 3.18 Seismogram of Suusamyr aftershock showing Pn and Pg phases.
- 3.19a Deep aftershock showing clear S phase
- 3.20a Near-field phases observed at AAK
- 3.20b Near-field phases observed at AML
- 3.21 Example of near-field phases observed in Pasadena
- 3.22 Apparent permanent offset observed at AML.
- 3.23 Apparent permanent offset observed at AML showing displacement beginning with P arrival.
- 3.24 Plots of offset versus amplitude, magnitude and P-S time
- 3.25 Seismogram of nuclear explosion at Lop Nor test site.

- 3.26 Map of Hindu Kush earthquakes and CMT solutions in relationship to KNET network.
 - 3.29 Seismograms of deep Hindu Kush earthquakes
- Chapter 4 The 1992 Suusamy, Kyrgyzstan, earthquake
- 4.1 Map of the northern Tien Shan
 - 4.2 Map of the Suusamyr aftershock sequence and recording stations
 - 4.3 Map of selected locations recorded by KNET and the aftershock network.
 - 4.4 Map and cross sections of best located aftershocks.
 - 4.5 Maps of earthquakes at different depths.
 - 4.6 Focal mechanisms for the largest aftershocks
 - 4.7 Comparison of results from the teleseismic body wave inversion
 - 4.8 Demonstration of results with a multiple source
 - 4.9 Testing sensitivity of body-wave inversion
 - 4.10 Profile of aftershocks and topography

CHAPTER 1:

Introduction

Scattered all across Central Asia, from the dusty plains of Turkmenistan to the airy peaks of the Tien Shan, seismometers quietly record every bump and twitch of the earth. The first instruments to record and locate the earthquakes that strike this seismically active area were installed in the 1920s [Galperin *et al.*, 1986]. Since then, the sophistication of both seismic instrumentation and data analysis techniques has greatly increased. In this thesis I use seismic data to determine the structure of the crust and upper mantle under the Hindu Kush (Chapter 2), to examine the source parameters of a large and damaging earthquake (Chapter 4), as well as to determine earthquake locations (Chapters 3 and 4).

The purpose of these studies is twofold. First, to help understand the active tectonic processes in this complex and poorly understood area. Second, and more importantly, to aid in evaluating the potential seismic hazard in this region which has suffered many catastrophic earthquakes in the past.

In Chapter 2, I apply the technique of teleseismic arrival time tomography to study the deep structure of the Hindu Kush and Pamir seismic zones. These enigmatic and highly active seismic zones possess a dipping zone of earthquakes similar to many subduction zones, yet lack an obvious surface expression. Previous studies had indicated both high mantle velocities [e.g. Vinnik and Lukk , 1975] and low mantle velocities [Roecker *et al.*, 1982]. My study shows that the deep part of the Hindu Kush zone is closely associated with high mantle velocities. These anomalously high velocities suggest that part of the zone may be subducted oceanic lithosphere. These results help resolve the discrepancy between previous studies and put constraints on the tectonic history of the Hindu Kush seismic zone.

The third chapter describes a new seismic network (KNET) deployed around the city of Bishkek in northern Kyrgyzstan and presents an analysis of the local seismicity recorded by this network, with emphasis on the seismicity near the city of Bishkek. The

seismicity near Bishkek helps to define the possible seismogenic structures near this large city and shows the seismic style of an active thrust belt. Chapter 3 also shows examples of data recorded by KNET including near-field effects and anomalous regional seismograms from the Hindu Kush zone.

The fourth chapter studies the Suusamyr earthquake, a large (Ms 7.4) earthquake that struck northern Kyrgyzstan on August 19, 1992. This destructive earthquake was not only first well-recorded large event in the northern Tien Shan but is also one of the largest well-recorded continental thrust events in the world today. Consequently, the study of this event yields information about active faulting in the region, seismic hazard and the geometry of thrust faults.

References

- Galperin, E. I., I. L. Nersesov, and R.M. Galperina, *Borehole seismology and the study of the seismic regime of large industrial centers*, D. Reidel Publishing Co., 1986.
- Lukk, A. A., and L. P. Vinnik, Tectonic interpretation of the deep structure of the Pamirs Engl. Transl., *Geotectonics*, 5, 300-304 , 1975.
- Roecker, S. W., Velocity structure of the Pamir-Hindu Kush region: Possible evidence of subducted crust, *J. Geophys. Res.*,87, 945-959, 1982.

Figure 1.1 (top) Map showing collision-induced tectonics of Central Asian region. Arrows indicate relative movement between regions. After Thomas et al., [1993].

(bottom) Map of same area showing earthquakes of magnitude 5.5 and above since 1977. Light gray represent elevations above 2000 m, darker gray is elevations above 4000m. The 'Hindu Kush' rectangle refers to the area studied in Chapter 1 while 'KNET' refers to the area covered in Chapter 2 and 3.